

Research

NSAIDs, Antithrombotic Therapy After Myocardial Infarction 805

Use of nonsteroidal anti-inflammatory drugs (NSAIDs) is discouraged after myocardial infarction; however, use of NSAIDs is widespread. In an analysis of Danish health and mortality administrative registries, Olsen and colleagues examined the association of NSAID use with risk of bleeding and cardiovascular events in 61 971 patients with incident myocardial infarction. The authors found that among patients receiving antithrombotic therapy after myocardial infarction, concomitant use of NSAIDs was associated with increased risk of bleeding and excess thrombotic events during a median follow-up of 3.5 years. In an Editorial, Campbell and Moliterno discuss hazards of adding NSAIDs to antithrombotic therapy after myocardial infarction.

📖 Editorial 801

Gene Variant Associated With Vincristine-Related Neuropathy 815

Vincristine—a part of every treatment protocol for childhood acute lymphoblastic leukemia (ALL)—is often associated with a dose-limiting peripheral neuropathy. Diouf and colleagues performed a genome-wide single-nucleotide polymorphism analysis in a cohort of 321 patients receiving treatment for childhood ALL and found that a polymorphism in the promoter region of the *CEP72* gene was associated with increased risk and severity of vincristine-related peripheral neuropathy. In an Editorial, McLeod discusses benefits and risks associated with personalized medicine.

📖 Editorial 803

📺 Author Video Interview jama.com

Management of Oral Anticoagulant-Associated ICH 824

Data to inform the acute and long-term treatment of patients with oral anticoagulant-associated intracerebral hemorrhage (OAC-ICH) are limited. In a retrospective cohort study involving 1176 patients with OAC-ICH, Kuramatsu and colleagues assessed the associations of anticoagulation-reversal and blood pressure levels with hematoma enlargement, and the association of OAC-resumption with patient outcomes. The authors report that reduced rates of hematoma enlargement were associated with systolic blood pressure less than 160 mm Hg at 4 hours and reversal of international normalized ratio levels to less than 1.3 within 4 hours of admission. Resumption of anticoagulant therapy was associated with a lower risk of ischemic events without increased bleeding complications.



Humanities

The Art of JAMA
778 *By the River*, 1912.
Edward W. Redfield (1869-1965).

Poetry and Medicine
860 Compline

JAMA Revisited
861 The Effects of Weather on Pain, Comfort and Efficiency

Opinion

Viewpoint

791 Genomic Engineering and the Future of Medicine
JA Doudna

793 Sharing Clinical Trial Data: Maximizing Benefits, Minimizing Risk
B Lo

795 Optimizing High-Risk Care Management
BW Powers, SK Chaguturu, and TG Ferris

797 Medicare and Care Coordination: Expanding the Clinician's Toolbox
L Aronson, CA Bautista, and K Covinsky

A Piece of My Mind

799 For What It's Worth
CA Humikowski

Editorial

801 Potential Hazards of Adding Nonsteroidal Anti-inflammatory Drugs to Antithrombotic Therapy After Myocardial Infarction: Time for More Than a Gut Check
CL Campbell and DJ Moliterno

803 Precision Medicine to Improve the Risk and Benefit of Cancer Care: Genetic Factors in Vincristine-Related Neuropathy
HL McLeod

LETTERS

Research Letter

853 Trends in the Proportion of Patients With Lung Cancer Meeting Screening Criteria
Y Wang and Coauthors

Comment & Response

855 Assessing the Benefit of Vascular Closure Devices After Femoral Artery Puncture

856 Male Body Image and Weight-Related Disorders

857 Alternative Pricing Strategies for Cancer Drugs

858 Correction

Editor in Chief
Howard Bauchner, MD

131 YEARS
OF CONTINUOUS
PUBLICATION

Clinical Review & Education

GFR, Albuminuria for Detection and Staging of Kidney Disease 837

Laboratory measurements are required to detect asymptomatic and early-stage kidney disease. Levey and colleagues summarize evidence from recent guidelines and reviews supporting the use of glomerular filtration rate (GFR) and albuminuria for detection and staging of acute and chronic kidney disease in adults. The authors review clinical characteristics of acute and chronic kidney disease, provide guidance for initial and confirmatory assessment of GFR and albuminuria for diagnosis and staging, and discuss indications for testing in special populations and assessing kidney function for drug dosing.

Continuing Medical Education jamanetworkcme.com

Hydroxychloroquine and the Retina 847

Patients with rheumatoid arthritis, systemic lupus erythematosus, and other autoimmune diseases are often treated with hydroxychloroquine—placing them at risk of irreversible toxic maculopathy. An article in *JAMA Ophthalmology* reported that many patients are not undergoing recommended routine monitoring for this serious adverse effect. In this From the JAMA Network article, Marmor and Melles discuss factors to consider in hydroxychloroquine dosing and the importance of annual screening to detect early signs of retinal toxicity.

Elevated Lactate Levels in a Non-Critically Ill Patient 849

This JAMA Diagnostic Test Interpretation article by Chen and colleagues presents a man with diffuse B-cell lymphoma involving the abdomen who was seen in the emergency department after 5 days of nausea and vomiting. He was afebrile and normotensive and he had mild tachycardia. Physical examination revealed a large, firm mass near the umbilicus, without abdominal rigidity or rebound tenderness. He had an elevated serum lactate level and low bicarbonate, low $Paco_2$, and marginally acidotic pH venous blood gas results. How would you interpret these findings?



JAMA Patient Page
868 Advance Directives

NEWS & ANALYSIS

Medical News & Perspectives

782 Efforts Seek to Develop Systematic Ways to Objectively Assess Surgeons' Skills

784 On Fear, Distrust, and Ebola

785 Health Agencies Update Proposed Rule Seeks to Safeguard Against Phased-Out Chemicals

Officials Strive to Help Consumers Shop for Health Insurance Benefits

Electric Stimulation Device Approved to Treat Obesity

786 Clinical Trials Update
Glaucoma Drug Significantly Slows the Progression of Vision Loss

Trials Will Assess Effect of Flexible Hours for Trainees

Progesterone Offers No Benefits in Traumatic Brain Injury

787 News From the CDC
Toxic Puffer Fish Species Illegally Imported to United States

Cholesterol-Lowering Drugs Taken by 28% of US Population

Online @ jama.com



Editor's Audio Summary

Howard Bauchner, MD, summarizes and comments on this week's issue.

Author Reading



AUDIO Jennifer Doudna, MD, reads her Viewpoint "Genomic Engineering and the Future of Medicine."

Author Interview

VIDEO Interview with William E. Evans, PharmD, author of "Association of an Inherited Genetic Variant With Vincristine-Related Peripheral Neuropathy in Children With Acute Lymphoblastic Leukemia"

Departments

775 Staff Listing

851 CME Questions

862 JAMA Network Abstracts

864 Classified Advertising

865 Journal Advertiser Index

867 Contact Information

Instructions for Authors

jama.com/public/instructionsforauthors.aspx