In This Issue of JAMA

October 2, 2018 Volume 320, Number 13, Pages 1295-1396

Research

Physical Therapy vs Surgery for Meniscal Tears

1328

Opinion Viewpoint

AS Fauri

1315 Addressing the Tuberculosis

Epidemic: 21st Century Research

1317 Crossing the Global Health Care

Quality Chasm: A Key Component of Universal Health Coverage

D Berwick, M Snair, and S Nishtar

and Chronic Disease: Emerging

1321 Aging as a Biological Target

N Barzilai, AM Cuervo, and S Austad

1323 From Lifespan to Healthspan

1326 Physical Therapy or Arthroscopic

Surgery for Treatment of Meniscal

Lazrawi, HT Gold, and JD Zuckerman

Tears: Is Noninferiority Enough?

for Prevention and Therapy

1319 Aging, Cell Senescence,

Therapeutic Strategies

SJ Olstransky

Z5 Sager

Editorial

LETTERS

Research Letter

A Piece of My Mind 1325 Nurturing Medicine

T Tchkonia and JL Kirldand

for an Ancient Disease



Arthroscopic partial meniscectomy is among the most frequently performed procedures in orthopedic surgery, but some evidence suggests that physical therapy may have a similar long-term benefit. Van de Graaf and colleagues for the Escape Research Group randomized 321 patients with nonobstructive meniscal tears and found that physical therapy was noninferior to arthroscopic partial meniscectomy for improving patient-reported

knee function over 24-month follow-up. In an Editorial, Jazrawi and colleagues suggest that many patients with meniscal tears associated with degenerative osteoarthritis may respond to nonoperative treatment.

Editorial 1326

■ CME jamanetwork.com/learning

Racial Differences in Risk Factors for Hypertension

Understanding racial differences in the prevalence of risk factors for hypertension could guide efforts to reduce disparities in cardiovascular disease. Howard and colleagues followed 6897 black and white adults for a median of 9.4 years and found that Southern diet score, dietary ratio of sodium to potassium, and education level mediated racial differences in the incidence of hypertension.

☐ CME jamanetwork.com/learning

Outcomes of Treatment for Subclinical Hypothyroidism

It is unclear whether thyroid hormone therapy is beneficial for subclinical hypothyroidism. defined as elevated thyroid-stimulating hormone with normal free thyroxine. In a meta-analysis of 21 trials with 2192 adults who had subclinical hypothyroidism, Feller and colleagues found that the use of thyroid hormone therapy was not associated with improvements in general quality of life or thyroid-related symptoms.

1379 Electronic Cigarette Sales in the United States, 2013-2017 BA King and Coauthors

1380 Presence of High-Intensity Sweeteners in Popular Cigarillos of Varying Flavor Profiles **HC Erythropel and Coauthors**

Comment & Response

1383 Glyburide vs Insulin for Gestational Diabetes

1384 Acupuncture for Women Undergoing In Vitro Fertilization

1386 Factors Associated With Variation in Outcomes in Barlatric Surgery Centers of Excellence

1387 Correction

Humanities

Poetry and Medicine 1388 Vitreous 1 Karetnick

JAMA Revisited 1389 The Book Forum



Editor in Chief Howard Bauchner, MD

135 YEARS OF CONTINUOUS PHRI ICATION

iama.com

In This Issue of JAMA

October 2, 2018 Volume 320, Number 13 Pages 1295-1396

Clinical Review & Education

Treatment of Rheumatoid Arthritis



Control of rheumatoid arthritis can be achieved with disease-modifying antirheumatic drugs, which reduce inflammation, improve physical function, and slow the progression of joint damage. Aletaha and Smolen review the clinical management of patients using a treat-to-target strategy.

Evaluation of an Adrenal Incidentaloma

1373

1360

This JAMA Diagnostic Test Interpretation article by Ospina and colleagues presents a 66-year-old man with an adrenal mass discovered during radiological evaluation for a reason other than suspected adrenal disease. Laboratory results indicated possible autonomous production of cortisol. What would you do next?

Ciprofloxacin Otic Suspension for Acute Otitis Externa

1375

Topical antibiotic therapy is the treatment of choice for acute otitis externa, but a typical course of treatment is 7 to 10 days. This Medical Letter on Drugs and Therapeutics discusses the efficacy and safety of a single dose of ciprofloxacin for acute otitis externa caused by *Pseudomonas aeruginosa* or *Staphylococcus aureus*.





JAMA Patient Page 1396 Tinea Versicolor

NEWS & ANALYSIS

Medical News & Perspectives

1303 Cardiologist Trades Stern Cells for Cell-based Meat

The JAMA Forum

1306 Given Their Potential for Harm, It's Time to Focus on the Safety of Supplements

Bench to Bedside

1307 Gut Microbes May Account for the Antiseizure Effects of the Ketogenic Diet

1309 News From the FDA

Rare Infection Linked With SGLT-2 Inhibitor Use

Tretinoin in New Lotion Formula

New Drug Treats Rare, Debilitating Neurotrophic Keratitis

Online @ jama.com



Editor's Audio Summary

Phil B. Fontanarosa, MD, MBA summarizes and comments on this week's issue.

Author Audio Interview

Evan Wood, MD. PhD, author of "Will This Hospitalized Patient Develop Severe Alcohol Withdrawal Syndrome? The Rational Clinical Examination Systematic Review"

Audio Interview

Uma Valeti, MD, Memphis Meats, profiled in "Cardiologist Trades Stem Cells for Cell-Based Meat"

Visual Abstract

"Effect of Early Surgery vs Physical Therapy on Knee Function Among Patients With Nonobstructive Meniscal Tears: The ESCAPE Randomized Clinical Trial" Departments

1301 Staff Listing

1377 CME Questions

1390 Classified Advertising

1393 Journal Advertiser Index

1395 Contact Information

Instructions for Authors

jama.com/public /instructionsforauthors.aspx